arXiv:1112.5442v1 [math.DG] 22 Dec 2011

A distinguished Riemannian geometrization for quadratic Hamiltonians of polymomenta

Alexandru Oană and Mircea Neagu

Abstract

In this paper we construct a distinguished Riemannian geometrization on the dual 1-jet space $J^{1*}(\mathcal{T}, M)$ for the multi-time quadratic Hamiltonian function

$$H = h_{ab}(t)g^{ij}(t,x)p_i^a p_j^b + U_{(a)}^{(i)}(t,x)p_i^a + \mathcal{F}(t,x).$$

Our geometrization includes a nonlinear connection N, a generalized Cartan canonical N-linear connection $C\Gamma(N)$ (together with its local d-torsions and d-curvatures), naturally provided by the given quadratic Hamiltonian function depending on polymomenta.

2010 Mathematics Subject Classification: 70S05, 53C07, 53C80.

Key words and phrases: dual 1-jet spaces, metrical multi-time Hamilton spaces, nonlinear connections, generalized Cartan canonical N-linear connection, d-torsions and d-curvatures.

1 Short introduction

In the last decades, numerous scientists were preoccupied by the geometrization of Hamiltonians depending on polymomenta. In such a perspective, we point out that the Hamiltonian geometrizations are achieved in three distinct ways:

- the *multisymplectic Hamiltonian geometry* − developed by Gotay, Isenberg, Marsden, Montgomery and their co-workers (see [10], [9]);
- the *polysymplectic Hamiltonian geometry* − elaborated by Giachetta, Mangiarotti and Sardanashvily (see [7], [8]);
- the *De Donder-Weyl Hamiltonian geometry* − studied by Kanatchikov (see [11], [12], [13]).

In such a geometrical context, the recent studies of Atanasiu and Neagu (see the papers [4], [5] and [6]) are initiating the new way of distinguished Riemannian geometrization for Hamiltonians depending on polymomenta, which is in fact a natural "multi-time" extension of the already classical Hamiltonian geometry on cotangent bundles synthesized in the Miron et al.'s book [16]. Note that our distinguished Riemannian geometrization for Hamiltonians depending on polymomenta is different one by all three Hamiltonian geometrizations from above (multisymplectic, polysymplectic and De Donder-Weyl).

2 Metrical multi-time Hamilton spaces

Let us consider that $h = (h_{ab}(t))$ is a semi-Riemannian metric on the "multi-time" (*temporal*) manifold $\mathcal{T}^m$, where $m = \dim \mathcal{T}$. Let $g = \left(g^{ij}(t^c, x^k, p^c_k)\right)$ be a symmetric d-tensor on the dual 1-jet space $E^* = J^{1*}(\mathcal{T}, M^n)$, which has the rank $n = \dim M$ and a constant signature. At the same time, let us consider a smooth multi-time Hamiltonian function

$$E^* \ni (t^a, x^i, p^a_i) \to H(t^a, x^i, p^a_i) \in \mathbb{R},$$

which yields the *fundamental vertical metrical d-tensor*

$$G^{(i)(j)}_{(a)(b)} = \frac{1}{2}\frac{\partial^2 H}{\partial p^a_i \partial p^b_j},$$

where $a, b = 1, .., m$ and $i, j = 1, ..., n$.

Definition 1 *A multi-time Hamiltonian function $H : E^* \to \mathbb{R}$, having the fundamental vertical metrical d-tensor of the form*

$$G^{(i)(j)}_{(a)(b)}(t^c, x^k, p^c_k) = \frac{1}{2}\frac{\partial^2 H}{\partial p^a_i \partial p^b_j} = h_{ab}(t^c) g^{ij}(t^c, x^k, p^c_k),$$

is called a ***Kronecker h-regular multi-time Hamiltonian function****.*

In such a context, we can introduce the following important geometrical concept:

Definition 2 *A pair $MH^n_m = (E^* = J^{1*}(\mathcal{T}, M), H)$, where $m = \dim \mathcal{T}$ and $n = \dim M$, consisting of the dual 1-jet space and a Kronecker h-regular multi-time Hamiltonian function $H : E^* \to \mathbb{R}$, is called a* ***multi-time Hamilton space****.*

Remark 3 *In the particular case $(\mathcal{T}, h) = (\mathbb{R}, \delta)$, a* ***"single-time" Hamilton space*** *will be also called a* ***relativistic rheonomic Hamilton space*** *and it will be denoted by $RRH^n = (J^{1*}(\mathbb{R}, M), H)$.*

Example 4 *Let us consider the Kronecker h-regular multi-time Hamiltonian function $H_1 : E^* \to \mathbb{R}$ given by*

$$H_1 = \frac{1}{mc} h_{ab}(t) \varphi^{ij}(x) p^a_i p^b_j, \tag{1}$$

where $h_{ab}(t)$ ($\varphi_{ij}(x)$, respectively) is a semi-Riemannian metric on the temporal (spatial, respectively) manifold $\mathcal{T}$ (M, respectively) having the physical meaning of ***gravitational potentials****, and m and c are the known constants from*

Theoretical Physics representing the ***mass of the test body*** *and the* ***speed of light****. Then, the multi-time Hamilton space* $\mathcal{G}MH^n_m = (E^*, H_1)$ *is called the* ***multi-time Hamilton space of the gravitational field****.*

Example 5 *If we consider on* E^* *a symmetric d-tensor field* $g^{ij}(t,x)$*, having the rank* n *and a constant signature, we can define the Kronecker h-regular multi-time Hamiltonian function* $H_2 : E^* \to \mathbb{R}$*, by setting*

$$H_2 = h_{ab}(t)g^{ij}(t,x)p^a_i p^b_j + U^{(i)}_{(a)}(t,x)p^a_i + \mathcal{F}(t,x), \tag{2}$$

where $U^{(i)}_{(a)}(t,x)$ *is a d-tensor field on* E^**, and* $\mathcal{F}(t,x)$ *is a function on* E^**. Then, the multi-time Hamilton space* $\mathcal{NEDM}H^n_m = (E^*, H_2)$ *is called the* ***non-autonomous multi-time Hamilton space of electrodynamics****. The dynamical character of the gravitational potentials* $g_{ij}(t,x)$ *(i.e., the dependence on the temporal coordinates* t^c*) motivated us to use the word* ***"non-autonomous".***

An important role for the subsequent development of our distinguished Riemannian geometrical theory for multi-time Hamilton spaces is represented by the following result (proved in the paper [4]):

Theorem 6 *If we have* $m = \dim \mathcal{T} \geq 2$*, then the following statements are equivalent:*

(i) *H is a Kronecker h-regular multi-time Hamiltonian function on* E^**.*

(ii) *The multi-time Hamiltonian function* H *reduces to a multi-time Hamiltonian function of non-autonomous electrodynamic type. In other words we have*

$$H = h_{ab}(t)g^{ij}(t,x)p^a_i p^b_j + U^{(i)}_{(a)}(t,x)p^a_i + \mathcal{F}(t,x). \tag{3}$$

Corollary 7 *The fundamental vertical metrical d-tensor of a Kronecker h-regular multi-time Hamiltonian function* H *has the form*

$$G^{(i)(j)}_{(a)(b)} = \frac{1}{2}\frac{\partial^2 H}{\partial p^a_i \partial p^b_j} = \begin{cases} h_{11}(t)g^{ij}(t,x^k,p^1_k), & m = \dim \mathcal{T} = 1 \\ h_{ab}(t^c)g^{ij}(t^c,x^k), & m = \dim \mathcal{T} \geq 2. \end{cases} \tag{4}$$

We recall that the transformations of coordinates on the dual 1-jet vector bundle $J^{1*}(\mathcal{T},M)$ are given by

$$\widetilde{t}^a = \widetilde{t}^a\left(t^b\right), \quad \widetilde{x}^i = \widetilde{x}^i\left(x^j\right), \quad \widetilde{p}^a_i = \frac{\partial x^j}{\partial \widetilde{x}^i}\frac{\partial \widetilde{t}^a}{\partial t^b}p^b_j,$$

where $\det\left(\partial\widetilde{t}^a/\partial t^b\right) \neq 0$ and $\det\left(\partial\widetilde{x}^i/\partial x^j\right) \neq 0$. In this context, let us introduce the following important geometrical concept:

Definition 8 *A pair of local functions on* $E^* = J^{1*}(\mathcal{T},M)$*, denoted by*

$$N = \left(\underset{1}{N}{}^{(a)}_{(i)b},\ \underset{2}{N}{}^{(a)}_{(i)j}\right),$$

whose local components obey the transformation rules

$$\begin{aligned}\widetilde{\underset{1}{N}}{}^{(b)}_{(j)c}\frac{\partial \widetilde{t}^c}{\partial t^a} &= \underset{1}{N}{}^{(c)}_{(k)a}\frac{\partial \widetilde{t}^b}{\partial t^c}\frac{\partial x^k}{\partial \widetilde{x}^j} - \frac{\partial \widetilde{p}^b_j}{\partial t^a},\\ \widetilde{\underset{2}{N}}{}^{(b)}_{(j)k}\frac{\partial \widetilde{x}^k}{\partial x^i} &= \underset{2}{N}{}^{(c)}_{(k)i}\frac{\partial \widetilde{t}^b}{\partial t^c}\frac{\partial x^k}{\partial \widetilde{x}^j} - \frac{\partial \widetilde{p}^b_j}{\partial x^i},\end{aligned}$$

is called a ***nonlinear connection*** *on E^*. The components $\underset{1}{N}{}^{(a)}_{(i)b}$ (resp. $\underset{2}{N}{}^{(a)}_{(i)j}$) are called the* ***temporal*** *(resp.* ***spatial****)* ***components*** *of N.*

Following now the geometrical ideas of Miron from [14], the paper [4] proves that any Kronecker h-regular multi-time Hamiltonian function H produces a natural nonlinear connection on the dual 1-jet space E^*, which depends only by the given Hamiltonian function H:

Theorem 9 *The pair of local functions $N = \left(\underset{1}{N}{}^{(a)}_{(i)b}, \underset{2}{N}{}^{(a)}_{(i)j}\right)$ on E^*, where (χ^a_{bc} are the Christoffel symbols of the semi-Riemannian temporal metric h_{ab})*

$$\begin{aligned}\underset{1}{N}{}^{(a)}_{(i)b} &= \chi^a_{bc}p^c_i,\\ \underset{2}{N}{}^{(a)}_{(i)j} &= \frac{h^{ab}}{4}\left[\frac{\partial g_{ij}}{\partial x^k}\frac{\partial H}{\partial p^b_k} - \frac{\partial g_{ij}}{\partial p^b_k}\frac{\partial H}{\partial x^k} + g_{ik}\frac{\partial^2 H}{\partial x^j \partial p^b_k} + g_{jk}\frac{\partial^2 H}{\partial x^i \partial p^b_k}\right],\end{aligned}$$

represents a nonlinear connection on E^*, which is called the ***canonical nonlinear connection of the multi-time Hamilton space*** *$MH^n_m = (E^*, H)$.*

Taking into account the Theorem 6 and using the *generalized spatial Christoffel symbols* of the d-tensor g_{ij}, which are given by

$$\Gamma^k_{ij} = \frac{g^{kl}}{2}\left(\frac{\partial g_{li}}{\partial x^j} + \frac{\partial g_{lj}}{\partial x^i} - \frac{\partial g_{ij}}{\partial x^l}\right),$$

we immediately obtain the following geometrical result:

Corollary 10 *For $m = \dim \mathcal{T} \geq 2$, the canonical nonlinear connection N of a multi-time Hamilton space $MH^n_m = (E^*, H)$, whose Hamiltonian function is given by (3), has the components*

$$\begin{aligned}\underset{1}{N}{}^{(a)}_{(i)b} &= \chi^a_{bc}p^c_i,\\ \underset{2}{N}{}^{(a)}_{(i)j} &= -\Gamma^k_{ij}p^a_k + T^{(a)}_{(i)j},\end{aligned}$$

where

$$T^{(a)}_{(i)j} = \frac{h^{ab}}{4}\left(U_{ib\bullet j} + U_{jb\bullet i}\right), \tag{5}$$

and

$$U_{ib} = g_{ik}U^{(k)}_{(b)}, \qquad U_{kb\bullet r} = \frac{\partial U_{kb}}{\partial x^r} - U_{sb}\Gamma^s_{kr}.$$

3 The Cartan canonical connection $C\Gamma(N)$ of a metrical multi-time Hamilton space

Let us consider that $MH^n_m = (J^{1*}(\mathcal{T}, M), H)$ is a multi-time Hamilton space, whose fundamental vertical metrical d-tensor is given by (4). Let

$$N = \left(\underset{1}{N}{}^{(a)}_{(i)b},\ \underset{2}{N}{}^{(a)}_{(i)j}\right)$$

be the canonical nonlinear connection of the multi-time Hamilton space MH^n_m.

Theorem 11 (the generalized Cartan canonical N-linear connection) *On the multi-time Hamilton space $MH^n_m = (J^{1*}(\mathcal{T}, M), H)$, endowed with the canonical nonlinear connection N, there exists an unique h-normal N-linear connection*

$$C\Gamma(N) = \left(\chi^a_{bc},\ A^i_{jc},\ H^i_{jk},\ C^{i(k)}_{j(c)}\right),$$

having the metrical properties:

(i) $g_{ij|k} = 0, \quad g^{ij}|^{(k)}_{(c)} = 0,$

(ii) $A^i_{jc} = \dfrac{g^{il}}{2}\dfrac{\delta g_{lj}}{\delta t^c}, \quad H^i_{jk} = H^i_{kj}, \quad C^{i(k)}_{j(c)} = C^{k(i)}_{j(c)},$

where "$_{/a}$", "$_{|k}$" and "$|^{(k)}_{(c)}$" represent the local covariant derivatives of the h-normal N-linear connection $C\Gamma(N)$.

Proof. Let $C\Gamma(N) = \left(\chi^a_{bc},\ A^i_{jc},\ H^i_{jk},\ C^{i(k)}_{j(c)}\right)$ be an h-normal N-linear connection, whose local coefficients are defined by the relations

$$\begin{aligned}
A^a_{bc} &= \chi^a_{bc}, \qquad A^i_{jc} = \frac{g^{il}}{2}\frac{\delta g_{lj}}{\delta t^c},\\
H^i_{jk} &= \frac{g^{ir}}{2}\left(\frac{\delta g_{jr}}{\delta x^k} + \frac{\delta g_{kr}}{\delta x^j} - \frac{\delta g_{jk}}{\delta x^r}\right),\\
C^{j(k)}_{i(c)} &= -\frac{g_{ir}}{2}\left(\frac{\partial g^{jr}}{\partial p^c_k} + \frac{\partial g^{kr}}{\partial p^c_j} - \frac{\partial g^{jk}}{\partial p^c_r}\right).
\end{aligned}$$

Taking into account the local expressions of the local covariant derivatives induced by the h-normal N-linear connection $C\Gamma(N)$, by local calculations, we deduce that $C\Gamma(N)$ satisfies conditions (i) and (ii).

Conversely, let us consider an h-normal N-linear connection

$$\tilde{C}\Gamma(N) = \left(\tilde{A}^a_{bc},\ \tilde{A}^i_{jc},\ \tilde{H}^i_{jk},\ \tilde{C}^{i(k)}_{j(c)}\right)$$

which satisfies conditions (i) and (ii). It follows that we have

$$\tilde{A}^a_{bc} = \chi^a_{bc}, \quad \tilde{A}^i_{jc} = \frac{g^{il}}{2}\frac{\delta g_{lj}}{\delta t^c}.$$

Moreover, the metrical condition $g_{ij|k} = 0$ is equivalent with

$$\frac{\delta g_{ij}}{\delta x^k} = g_{rj}\tilde{H}^r_{ik} + g_{ir}\tilde{H}^r_{jk}.$$

Applying now a Christoffel process to indices $\{i, j, k\}$, we find

$$\tilde{H}^i_{jk} = \frac{g^{ir}}{2}\left(\frac{\delta g_{jr}}{\delta x^k} + \frac{\delta g_{kr}}{\delta x^j} - \frac{\delta g_{jk}}{\delta x^r}\right).$$

By analogy, using the relations $C^{i(k)}_{j(c)} = C^{k(i)}_{j(c)}$ and $g^{ij}|^{(k)}_{(c)} = 0$, together with a Christoffel process applied to indices $\{i, j, k\}$, we obtain

$$\widetilde{C}^{j(k)}_{i(c)} = -\frac{g_{ir}}{2}\left(\frac{\partial g^{jr}}{\partial p^c_k} + \frac{\partial g^{kr}}{\partial p^c_j} - \frac{\partial g^{jk}}{\partial p^c_r}\right).$$

In conclusion, the uniqueness of the *generalized Cartan canonical connection* $C\Gamma(N)$ on the dual 1-jet space $E^* = J^{1*}(\mathcal{T}, M)$ is clear. ■

Remark 12 (i) *Replacing the canonical nonlinear connection* N *of the multi-time Hamilton space* MH^n_m *with an arbitrary nonlinear connection* $\hat{N}$*, the preceding Theorem holds good.*

(ii) *The generalized Cartan canonical connection* $C\Gamma(N)$ *of the multi-time Hamilton space* MH^n_m *verifies also the metrical properties*

$$h_{ab/c} = h_{ab|k} = h_{ab}|^{(k)}_{(c)} = 0, \quad g_{ij/c} = 0.$$

(iii) *In the case* $m = \dim\mathcal{T} \geq 2$*, the coefficients of the generalized Cartan canonical connection* $C\Gamma(N)$ *of the multi-time Hamilton space* MH^n_m *reduce to*

$$A^a_{bc} = \chi^a_{bc}, \quad A^i_{jc} = \frac{g^{il}}{2}\frac{\partial g_{lj}}{\partial t^c}, \quad H^i_{jk} = \Gamma^i_{jk}, \quad C^{i(k)}_{j(c)} = 0. \tag{6}$$

4 Local d-torsions and d-curvatures of the generalized Cartan canonical connection $C\Gamma(N)$

Applying the formulas that determine the local d-torsions and d-curvatures of an h-normal N-linear connection $D\Gamma(N)$ (see these formulas in [22]) to the generalized Cartan canonical connection $C\Gamma(N)$, we obtain the following important geometrical results:

Theorem 13 *The torsion tensor* $\mathbb{T}$ *of the generalized Cartan canonical connection* $C\Gamma(N)$ *of the multi-time Hamilton space* MH^n_m *is determined by the local* *d-components*

	$h_{\mathcal{T}}$	h_M		v	
	$m\geq 1$	$m=1$	$m\geq 2$	$m=1$	$m\geq 2$
$h_{\mathcal{T}}h_{\mathcal{T}}$	0	0	0	0	$R^{(f)}_{(r)ab}$
$h_M h_{\mathcal{T}}$	0	T^r_{1j}	T^r_{aj}	$R^{(1)}_{(r)1j}$	$R^{(f)}_{(r)aj}$
$vh_{\mathcal{T}}$	0	0	0	$P^{(1)\ (j)}_{(r)1(1)}$	$P^{(f)\ (j)}_{(r)a(b)}$
$h_M h_M$	0	0	0	$R^{(1)}_{(r)ij}$	$R^{(f)}_{(r)ij}$
vh_M	0	$P^{r(j)}_{i(1)}$	0	$P^{(1)\ (j)}_{(r)i(1)}$	0
vv	0	0	0	0	0

where

(i) *for* $m=\dim\mathcal{T}=1$, *we have*

$$T^r_{1j}=-A^r_{j1},\quad P^{r(j)}_{i(1)}=C^{r(j)}_{i(1)},\quad P^{(1)\ (j)}_{(r)1(1)}=\frac{\partial \underset{1}{N}{}^{(1)}_{(r)1}}{\partial p^1_j}+A^j_{r1}-\delta^j_r\chi^1_{11},$$

$$P^{(1)\ (j)}_{(r)i(1)}=\frac{\partial \underset{2}{N}{}^{(1)}_{(r)i}}{\partial p^1_j}+H^j_{ri},\quad R^{(1)}_{(r)1j}=\frac{\delta \underset{1}{N}{}^{(1)}_{(r)1}}{\delta x^j}-\frac{\delta \underset{2}{N}{}^{(1)}_{(r)j}}{\delta t},$$

$$R^{(1)}_{(r)ij}=\frac{\delta \underset{2}{N}{}^{(1)}_{(r)i}}{\delta x^j}-\frac{\delta \underset{2}{N}{}^{(1)}_{(r)j}}{\delta x^i};$$

(ii) *for* $m=\dim\mathcal{T}\geq 2$, *using the equality (5) and the notations*

$$\chi^c_{fab}=\frac{\partial\chi^c_{fa}}{\partial t^b}-\frac{\partial\chi^c_{fb}}{\partial t^a}+\chi^d_{fa}\chi^c_{db}-\chi^d_{fb}\chi^c_{da},$$

$$\mathfrak{R}^r_{kij}=\frac{\partial\Gamma^r_{ki}}{\partial x^j}-\frac{\partial\Gamma^r_{kj}}{\partial x^i}+\Gamma^p_{ki}\Gamma^r_{pj}-\Gamma^p_{kj}\Gamma^r_{pi},$$

we have

$$T^r_{aj}=-A^r_{ja},\quad P^{(f)\ (j)}_{(r)a(b)}=\delta^f_b A^j_{ra},\quad R^{(f)}_{(r)ab}=\chi^f_{gab}p^g_r,$$

$$R^{(f)}_{(r)aj}=-\frac{\partial \underset{2}{N}{}^{(f)}_{(r)j}}{\partial t^a}-\chi^f_{ca}T^{(c)}_{(r)j},$$

$$R^{(f)}_{(r)ij}=-\mathfrak{R}^k_{rij}p^f_k+\left[T^{(f)}_{(r)i|j}-T^{(f)}_{(r)j|i}\right].$$

Theorem 14 *The curvature tensor* $\mathbb{R}$ *of the generalized Cartan canonical connection* $C\Gamma(N)$ *of the multi-time Hamilton space* MH^n_m *is determined by the* *following adapted local curvature d-tensors:*

	$h_{\mathcal{T}}$	h_M		v	
	$m \geq 1$	$m = 1$	$m \geq 2$	$m = 1$	$m \geq 2$
$h_{\mathcal{T}} h_{\mathcal{T}}$	χ^d_{abc}	0	R^l_{ibc}	0	$-R^{(d)(i)}_{(l)(a)bc}$
$h_M h_{\mathcal{T}}$	0	R^l_{i1k}	R^l_{ibk}	$-R^{(1)(l)}_{(i)(1)1k} = -R^l_{i1k}$	$-R^{(d)(i)}_{(l)(a)bk}$
$v h_{\mathcal{T}}$	0	$P^{l\ \ (k)}_{i1(1)}$	0	$-P^{(1)(l)\ (k)}_{(i)(1)1(1)} = -P^{l\ \ (k)}_{i1(1)}$	0
$h_M h_M$	0	R^l_{ijk}	$\mathfrak{R}^l_{ijk}$	$-R^{(1)(l)}_{(i)(1)jk} = -R^l_{ijk}$	$-R^{(d)(i)}_{(l)(a)jk}$
$v h_M$	0	$P^{l\ (k)}_{ij(1)}$	0	$-P^{(1)(l)\ (k)}_{(i)(1)j(1)} = -P^{l\ (k)}_{ij(1)}$	0
vv	0	$S^{l(j)(k)}_{i(1)(1)}$	0	$-S^{(1)(l)(j)(k)}_{(i)(1)(1)(1)} = -S^{l(j)(k)}_{i(1)(1)}$	0

where, for $m \geq 2$, we have the relations

$$-R^{(d)(i)}_{(l)(a)bc} = \delta^i_l \chi^d_{abc} - \delta^d_a R^i_{lbc}, \quad -R^{(d)(i)}_{(l)(a)bk} = -\delta^d_a R^i_{lbk}, \quad -R^{(d)(l)}_{(i)(a)jk} = -\delta^d_a \mathfrak{R}^l_{ijk},$$

and, generally, the following formulas are true:

(i) *for $m = \dim \mathcal{T} = 1$, we have $\chi^1_{111} = 0$ and*

$$R^l_{i1k} = \frac{\delta A^l_{i1}}{\delta x^k} - \frac{\delta H^l_{ik}}{\delta t} + A^r_{i1} H^l_{rk} - H^r_{ik} A^l_{r1} + C^{l(r)}_{i(1)} R^{(1)}_{(r)1k},$$

$$R^l_{ijk} = \frac{\delta H^l_{ij}}{\delta x^k} - \frac{\delta H^l_{ik}}{\delta x^j} + H^r_{ij} H^l_{rk} - H^r_{ik} H^l_{rj} + C^{l(r)}_{i(1)} R^{(1)}_{(r)jk},$$

$$P^{l\ \ (k)}_{i1(1)} = \frac{\partial A^l_{i1}}{\partial p^1_k} - C^{l(k)}_{i(1)/1} + C^{l(r)}_{i(1)} P^{(1)\ \ (k)}_{(r)1(1)},$$

$$P^{l\ (k)}_{ij(1)} = \frac{\partial H^l_{ij}}{\partial p^1_k} - C^{l(k)}_{i(1)|j} + C^{l(r)}_{i(1)} P^{(1)\ \ (k)}_{(r)j(1)},$$

$$S^{l(j)(k)}_{i(1)(1)} = \frac{\partial C^{l(j)}_{i(1)}}{\partial p^1_k} - \frac{\partial C^{l(k)}_{i(1)}}{\partial p^1_j} + C^{r(j)}_{i(1)} C^{l(k)}_{r(1)} - C^{r(k)}_{i(1)} C^{l(j)}_{r(1)};$$

(ii) *for $m = \dim \mathcal{T} \geq 2$, we have*

$$\chi^d_{abc} = \frac{\partial \chi^d_{ab}}{\partial t^c} - \frac{\partial \chi^d_{ac}}{\partial t^b} + \chi^f_{ab} \chi^d_{fc} - \chi^f_{ac} \chi^d_{fb},$$

$$R^l_{ibc} = \frac{\partial A^l_{ib}}{\partial t^c} - \frac{\partial A^l_{ic}}{\partial t^b} + A^r_{ib} A^l_{rc} - A^r_{ic} A^l_{rb},$$

$$R^l_{ibk} = \frac{\partial A^l_{ib}}{\partial x^k} - \frac{\partial \Gamma^l_{ik}}{\partial t^b} + A^r_{ib} \Gamma^l_{rk} - \Gamma^r_{ik} A^l_{rb},$$

$$\mathfrak{R}^l_{ijk} = \frac{\partial \Gamma^l_{ij}}{\partial x^k} - \frac{\partial \Gamma^l_{ik}}{\partial x^j} + \Gamma^r_{ij} \Gamma^l_{rk} - \Gamma^r_{ik} \Gamma^l_{rj}.$$

Acknowledgements. The authors of this paper thank to Professor Gh. Atanasiu for our interesting and useful discussions on this research topic.

References

[1] G.S. Asanov, *Jet extension of Finslerian gauge approach*, Fortschritte der Physik, vol. **38**, no. **8** (1990), 571-610.

[2] Gh. Atanasiu, *The invariant expression of Hamilton geometry*, Tensor N.S., vol. **47**, no. **3** (1988), 225-234.

[3] Gh. Atanasiu, F.C. Klepp, *Nonlinear connections in cotangent bundle*, Publ. Math. Debrecen, Hungary, vol. **39**, no. **1-2** (1991), 107-111.

[4] Gh. Atanasiu, M. Neagu, *Canonical nonlinear connections in the multi-time Hamilton geometry*, Balkan Journal of Geometry and Its Applications, vol. **14**, no. **2** (2009), 1-12.

[5] Gh. Atanasiu, M. Neagu, *Distinguished tensors and Poisson brackets in the multi-time Hamilton geometry*, BSG Proceedings **16**, Geometry Balkan Press, Bucharest (2009), 12-27.

[6] Gh. Atanasiu, M. Neagu, *Distinguished torsion, curvature and deflection tensors in the multi-time Hamilton geometry*, Electronic Journal "Differential Geometry-Dynamical Systems", vol. **11** (2009), 20-40.

[7] G. Giachetta, L. Mangiarotti, G. Sardanashvily, *Covariant Hamiltonian field theory*, http://arxiv.org/hep-th/9904062 (1999).

[8] G. Giachetta, L. Mangiarotti, G. Sardanashvily, *Polysymplectic Hamiltonian formalism and some quantum outcomes*, http://arxiv.org/hep-th/0411005 (2004).

[9] M. Gotay, J. Isenberg, J.E. Marsden, R. Montgomery, *Momentum maps and classical fields. Part I. Covariant field theory*, http://arxiv.org/physics/9801019 (2004).

[10] M. Gotay, J. Isenberg, J.E. Marsden, *Momentum maps and classical fields. Part II. Canonical analysis of field theories*, http://arxiv.org/math-ph/0411032 (2004).

[11] I.V. Kanatchikov, *Basic structures of the covariant canonical formalism for fields based on the De Donder-Weyl theory*, http://arxiv.org/hep-th/9410238 (1994).

[12] I.V. Kanatchikov, *On quantization of field theories in polymomentum variables*, AIP Conf. Proc., vol. **453**, Issue **1** (1998), 356-367.

[13] I.V. Kanatchikov, *On the canonical structure of the De Donder-Weyl covariant Hamiltonian formulation of field theory I. Graded Poisson brackets and equations of motion*, http://arxiv.org/hep-th/9312162 (1993).

[14] R. Miron, *Hamilton geometry*, An. Şt. "Al. I. Cuza" Univ., Iaşi, Romania, vol. **35** (1989), 33-67.

[15] R. Miron, M. Anastasiei, *The Geometry of Lagrange Spaces: Theory and Applications*, Kluwer Academic Publishers, 1994.

[16] R. Miron, D. Hrimiuc, H. Shimada, S.V. Sabău, *The Geometry of Hamilton and Lagrange Spaces*, Kluwer Academic Publishers, 2001.

[17] R. Miron, M.S. Kirkovits, M. Anastasiei, *A geometrical model for variational problems of multiple integrals*, Proc. Conf. Diff. Geom. and Appl., Dubrovnik, Yugoslavia, June 26-July 3, (1988), 8-25.

[18] M. Neagu, *Ricci and Bianchi identities for h-normal Γ-linear connections on $J^1(T, M)$*, Hindawi Publishing Corporation, International Journal of Mathematics and Mathematical Sciences, no. **34** (2003), 2177-2192.

[19] M. Neagu, *Riemann-Lagrange Geometry on* 1*-Jet Spaces*, Matrix Rom, Bucharest, 2005.

[20] M. Neagu, C. Udrişte, *Torsion, curvature and deflection d-tensors on $J^1(T, M)$*, Balkan Journal of Geometry and Its Applications, vol. **6**, no. **1** (2001), 29-45.

[21] M. Neagu, C. Udrişte, A. Oană, *Multi-time dependent sprays and h-traceless maps,* Balkan Journal of Geometry and Its Applications, vol. **10**, no. **2** (2005), 76-92.

[22] A. Oană, M. Neagu, *The local description of the Ricci and Bianchi identities for an h-normal N-linear connection on the dual* 1*-jet space $J^{1*}(\mathcal{T}, M)$,* http://arxiv.org/math.DG/1111.4173 (2011).

[23] P.J. Olver, *Applications of Lie Groups to Differential Equations*, Graduate Texts in Mathematics, vol. **107**, Springer-Verlag, New York, 1986.

[24] D.J. Saunders, *The Geometry of Jet Bundles*, Cambridge University Press, New York, London, 1989.

Alexandru OANĂ and Mircea NEAGU
University Transilvania of Braşov,
Department of Mathematics - Informatics,
Blvd. Iuliu Maniu, no. 50, Braşov 500091, Romania.
E-mails: alexandru.oana@unitbv.ro, mircea.neagu@unitbv.ro